



II Congreso

Sociedad Hispanoamericana de Hernia

3, 4 y 5 de mayo de 2018

Aula Magna de la Fundación Jiménez Díaz

Madrid - España



CIRUGIA MINI INVASIVA EN HERNIA INGUINAL... Por qué ?

Luis Gabriel González Higuera
Hospital Central Policía Nacional
Bogotá-Colombia



Recurrences After Conventional Anterior and Laparoscopic Inguinal Hernia Repair

A Randomized Comparison

Mike S. L. Liem, MD,* Eino B. van Duyn, MD,* Yolanda van der Graaf, MD,† and Theo J. M. V. van Vroonhoven, MD* on behalf of the Coala Trial Group

Menor recidiva



Menor dolor POP



Compara técnicas con y sin tensión



Seguimiento a 5 años



Meta-analysis of randomized clinical trials comparing open and laparoscopic inguinal hernia repair

M. A. Memon¹, N. J. Cooper², B. Memon³, M. I. Memon⁴ and K. R. Abrams²

¹Department of Surgery, Nottingham City Hospital, Nottingham, ²Department of Epidemiology and Public Health, University of Leicester, Leicester,

³Bolton Institute, Bolton and ⁴Department of Community Health, Preston Primary Care Trust and Multicultural Studies Centre, Bolton Institute, Bolton, UK

Correspondence to: Mr M. A. Memon, Astley House, Whitehall Road, Darwen BB3 2LH, UK (e-mail: mmemon@yahoo.com)

Alta precoz.

Retorno rápido actividades diarias.

Menos complicaciones.

Mayor tiempo operatorio.

Mínimo aumento de la recurrencia



From *The Cochrane Library*, Issue 3, 2004. Chichester, UK: John Wiley & Sons, Ltd. All rights reserved.

Laparoscopic techniques versus open techniques for inguinal hernia repair (Cochrane Review)

McCormack K, Scott NW, Go PMNYH, Ross S, Grant AM on behalf of the EU Hernia Trialists Collaboration

Mayor tiempo
quirúrgico

Ligero aumento
de
complicaciones
mayores

Menor
incapacidad
laboral

Rapido retorno a
ABC

Menor DICP

Menor frecuencia
de presentación
de parestesias y
alodinia

Menor recidiva
que la cirugía
abierta con y sin
uso de malla



Laparoscopic surgery for inguinal hernia repair: systematic review of effectiveness and economic evaluation

37 evaluación Qx y 14 evaluación económica.

Rápido retorno a actividades diarias.

Menos DICI y disestesias.

Menos complicaciones de la herida.

Menos seromas y hematomas.

Recidiva similar.

Mayor el costo directo...pero, si se considera la < incapacidad = > el costo de cx abierta



Laparoscopic surgery for inguinal hernia repair: systematic review of effectiveness and economic evaluation

UNILATERAL

- Mayor costo
- Mejor QOL

BILATERAL

- Mejor costo-efectividad con LAP.
- Mucho mejor en QOL

FUNDAMENTAL: Buen entrenamiento!



Randomized Clinical Trial of Total Extraperitoneal Inguinal Hernioplasty vs Lichtenstein Repair

A Long-term Follow-up Study

Hasan H. Eker, MD; Hester R. Langeveld, MD; Pieter J. Klitsie, MD; Martijne van't Riet, MD, PhD;
Laurents P. S. Stassen, MD, PhD; Wibbo F. Weidema, MD, PhD; Ewout W. Steyerberg, PhD;
Johan F. Lange, MD, PhD; Hendrik J. Bonjer, MD, PhD; Johannes Jeekel, MD, PhD

660 pacientes

5 años seguimiento.

Menor dolor crónico pop, ↓ parestesias (1x22%).

Rápido retorno al trabajo.

Mayor costo directo TEP, Costo total similar.

Mayor satisfacción de los pacientes con TEP.

Menor recurrencia con cirujanos expertos.



Laparoscopic Versus Open Inguinal Hernia Repair

Leandro Totti Cavazzola, MD, PhD^{a,*}, Michael J. Rosen, MD^b

KEY POINTS

- The laparoscopic approach to inguinal hernia surgery is a safe and reliable method with a similar recurrence rate as the open tension-free mesh repair.
- Because the laparoscopic approach shows clear advantages with regard to less chronic postoperative pain and numbness, fast return to normal activities, and a decrease in the incidence of wound infection and hematoma, it should be considered an appropriate approach for inguinal hernia surgery.
- The choice between the transabdominal preperitoneal (TAPP) procedure and the totally extraperitoneal (TEP) procedure should be based on patient and surgeon characteristics, because there is no evidence of superiority between either techniques.
- The use of lightweight mesh, with glue fixation (in TAPP procedures) or nonfixation (in TEP procedures), can achieve excellent results in the setting of a proficient surgeon.



Tailored approach in inguinal hernia repair – decision tree based on the guidelines

Ferdinand Köckerling* and Christine Schug-Pass

Department of Surgery, Centre for Minimally Invasive Surgery, Vivantes Hospital Berlin, Academic Teaching Hospital of Charité Medical School, Berlin, Germany

Edited by:

René H. Fortelny, Wilhelminenspital, Austria


Reviewed by:

*Frederik Helgstrand, University of Copenhagen, Denmark
Ulf Gunnarsson, Karolinska Institutet, Sweden*

***Correspondence:**

*Ferdinand Köckerling, Department of Surgery, Centre for Minimally Invasive Surgery, Vivantes Hospital Berlin, Academic Teaching Hospital of Charité Medical School, Neue Bergstrasse 6, Berlin 13585, Germany
e-mail: ferdinand.koeckerling@vivantes.de*


The endoscopic procedures TEP and TAPP and the open techniques Lichtenstein, Plug and Patch, and PHS currently represent the gold standard in inguinal hernia repair recommended in the guidelines of the European Hernia Society, the International Endohernia Society, and the European Association of Endoscopic Surgery. Eighty-two percent of experienced hernia surgeons use the "tailored approach," the differentiated use of the several inguinal hernia repair techniques depending on the findings of the patient, trying to minimize the risks. The following differential therapeutic situations must be distinguished in inguinal hernia repair: unilateral in men, unilateral in women, bilateral, scrotal, after previous pelvic and lower abdominal surgery, no general anesthesia possible, recurrence, and emergency surgery. Evidence-based guidelines and consensus conferences of experts give recommendations for the best approach in the individual situation of a patient. This review tries to summarize the recommendations of the various guidelines and to transfer them into a practical decision tree for the daily work of surgeons performing inguinal hernia repair.



CONTINUING MEDICAL EDUCATION

Evidence-Based Hernia Treatment in Adults

Dieter Berger



Head of Department
Prof. Dr. med. Dieter Berger



KLINIKUM
MITTELBADEN

Municipal Clinic Baden-Baden
Dep. of Visceral Surgery
Balger Strasse 50
D-76532 Baden-Baden
Secretary:
Mrs. Marion Weingart
Tel.: +49(0)7221/91-2351

Deutsches Ärzteblatt International
Dtsch Arztebl Int 2016; 113: 150

Accepted on 19 January
2016.

Hernia inguinal primaria

TABLE 1

Treatment options for primary inguinal hernia

	Conservative	Operative	open / anterior approach	laparoscopic / endoscopic
Unilateral hernia in a man, asymptomatic, non-progressive	+	+	+	+
Unilateral hernia in a man, symptomatic and/or progressive	–	+	+	+
Bilateral hernia in a man, asymptomatic, non-progressive	+	+	–	+
Bilateral hernia in a man, symptomatic and/or progressive	–	+	–	+
Hernia in a woman, unilateral/bilateral/asymptomatic/symptomatic/non-progressive/progressive	–	+	–	+

Hernia inguinal recurrente

TABLE 1

Treatment options for primary inguinal hernia

	Conserva- tive	Operative	open / anterior approach	laparoscopic / endoscopic
Unilateral hernia in a man, asymptomatic, non-progressive	+	+	+	+
Unilateral hernia in a man, symptomatic and/or progressive	–	+	+	+
Bilateral hernia in a man, asymptomatic, non-progressive	+	+	–	+
Bilateral hernia in a man, symptomatic and/or progressive	–	+	–	+
Hernia in a woman, unilateral/bilateral/asymptomatic/ symptomatic/non-progressive/progressive	–	+	–	+

Statements and Recommendations

<i>Statement</i>	When the surgeon has sufficient experience in the technique, laparo-endoscopic techniques do not take longer than Lichtenstein operations.	⊗⊗⊗□
<i>Statement</i>	When the surgeon has sufficient experience, no significant differences are observed in the perioperative complications needing reoperation between the laparo-endoscopic and Lichtenstein techniques.	⊗⊗⊗□
<i>Statement</i>	Laparo-endoscopic techniques have less chronic pain and a faster recovery than the Lichtenstein repair.	⊗⊗⊗□
<i>Statement</i>	The direct operative costs for laparo-endoscopic inguinal hernia repair are higher. That difference decreases when the total community costs are taken into account and the surgeon has sufficient experience.	⊗⊗⊗□
<i>Statement</i>	The learning curve for laparo-endoscopic techniques (especially TEP) is longer than for Lichtenstein. There are rare but severe complications mainly described early in the learning curve. It is imperative that laparo-endoscopic techniques be learned in a properly supervised manner in order to minimise complications..	⊗□□□
<i>Recommendation</i>	For male patients with primary unilateral inguinal hernia, a laparo-endoscopic technique is suggested because of a lower postoperative pain incidence and a reduction in chronic pain incidence, provided that a surgeon with specific expertise and sufficient resources is available. However, there are patient and hernia characteristics that warrant a Lichtenstein as first choice. (see chapter 7 on individualization)	⊗⊗⊗□ Weak downgraded

EXPERTICIA!!!



¿Por que Herniolap?

Menor DICP

Menor dolor
agudo.

Mejor evaluación
por los pacientes
(Q-alis).

Menor incapacidad
laboral.

Mas rápido ABC.

Menor tasa de
complicaciones de
la herida.

Se puede hacer
con costo mínimo
casi igual al de la
cx abierta.

A mayor difusión
de la técnica
mejores resultados
(<recidiva).

¡Porque Sí!

¡Gracias!





lugagonzalezh@hotmail.com

Especialista Cirugía General/Cirugía de Hernia
